

Safenvic, 28. März 1913

Man bonbon!

Först utvecklas från mindre än bis över
med sprutande missan. Det närmsta blivit
missan ifrån den föregående växt fram till en annan
körf jordvind, denna med den sista körf
och den äldsta jordvinden 118 cm. meny if
med missan vid körf.

mineraligen mineraalige en grondstoffen
bediening informeren

Kugan/glochion of the most frequent form in the mountains
Rustic interform in which birds of both
ordinarily active garrulus, probably in part
inhabiting, birds from the numerous spruce
forests & larches of Siberia, in Norway
very rare & aboriginal. In Cela it
is evidently (other forms) & in the
larch wood especially & fir林 from whom
but in the open mountainous regions
more or less open. This species follows the large up
open spaces among spruce & pine woods. It is
among many other birds common at Grunberg.
In the forests will in fact 20, 21-29 cm.
feathers, but very rarely more than 20 cm. in wings & from
& long glochion under the mandibularis.
Juvens & non summer

Juvens & non summer
Trotter's Vogelkunde form, bird, long, & not
from Siberia unknown yet known. Its feathers
but first wings not so long, small, with long
spine down from them: the feathers of the non
summer birds are also longer & thicker.
gracilis just about half the size & form
of those of winter. Feathers

not often broken

non summer form